

VU Research Portal

Strong Resemblance in the Amplitude of Oscillatory Brain Activity in Monozygotic Twins Is Not Caused by "Trivial" Similarities in the Composition of the Skull

van t Ent, D.; van Soelen, I.L.C.; Stam, C.J.; de Geus, E.J.C.; Boomsma, D.I.

published in

Human Brain Mapping

2009

DOI (link to publisher)

[10.1002/hbm.20656](https://doi.org/10.1002/hbm.20656)

[Link to publication in VU Research Portal](#)

citation for published version (APA)

van t Ent, D., van Soelen, I. L. C., Stam, C. J., de Geus, E. J. C., & Boomsma, D. I. (2009). Strong Resemblance in the Amplitude of Oscillatory Brain Activity in Monozygotic Twins Is Not Caused by "Trivial" Similarities in the Composition of the Skull. *Human Brain Mapping*, 30(7), 2142-2145. <https://doi.org/10.1002/hbm.20656>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

E-mail address:

vuresearchportal.ub@vu.nl

Strong Resemblance in the Amplitude of Oscillatory Brain Activity in Monozygotic Twins Is Not Caused by “Trivial” Similarities in the Composition of the Skull

Dennis van 't Ent,^{1,2*} Inge L.C. van Soelen,^{1,3} Cornelis J. Stam,⁴
Eco J.C. De Geus,¹ and Dorret I. Boomsma¹

¹Department of Biological Psychology, Vrije Universiteit, Amsterdam, The Netherlands

²Center for Neurogenomics and Cognitive Research (CNCR), VU University Medical Center, Amsterdam, The Netherlands

³Rudolf Magnus Institute of Neuroscience, Department of Psychiatry, University Medical Center Utrecht, The Netherlands

⁴Department of Clinical Neurophysiology and MEG Center, VU University Medical Center, Amsterdam, The Netherlands



Abstract: Previous twin studies have shown strong heritability of electroencephalogram amplitude characteristics, such as power spectra. However, it has been suggested that these high heritabilities may reflect “trivial” twin resemblance in intervening tissues such as the skull. Here we demonstrate strong monozygotic twin correlation ($0.79 < r < 0.88$) of eyes-closed resting-state magnetoencephalogram power, which is insensitive to intervening tissues. These results confirm that brain activity itself is highly heritable. *Hum Brain Mapp* 30:2142–2145, 2009. © 2008 Wiley-Liss, Inc.

Key words: brain electrical activity; magnetoencephalography; correlation study; monozygotic twins



INTRODUCTION

Electroencephalogram (EEG) measurements in humans consistently indicated a significant genetic influence on individual differences in electrical brain activity [van Beijsterveldt and Boomsma, 1994; Vogel and Motulsky, 1997]. For example, large-scale studies that compared similarities of EEG characteristics between genetically identical, mono-

zygotic (MZ) twins and nonidentical dizygotic (DZ) twins estimated high heritabilities for power in the classical EEG frequency bands. For power in delta, theta, alpha, and beta bands heritabilities range from 55% to 90% in children [Van Baal et al., 1996] and 70% to 90% in adolescents [Smit et al., 2006; van Beijsterveldt et al., 1996; Zietsch et al., 2007] and adults [Smit et al., 2005]. These findings suggest that, besides brain anatomy [Toga and Thompson, 2005], resting-state brain function is also under strong genetic control. However, EEG power is strongly influenced by electrical conductivity of tissue layers between the brain and scalp electrodes, in particular the skull [Babiloni et al., 1997; Leissner et al., 1970]. Heritability estimates of EEG power could be strongly inflated by greater MZ similarity for these intervening biological tissues which, as many other anatomical traits, are highly heritable themselves

*Correspondence to: Dennis van 't Ent, Department of Biological Psychology, Vrije Universiteit, van der Boechorststraat 1, 1081 BT, Amsterdam, The Netherlands. E-mail: d.vant.ent@psy.vu.nl

Received for publication 8 May 2008; Accepted 24 July 2008

DOI: 10.1002/hbm.20656

Published online 25 September 2008 in Wiley InterScience (www.interscience.wiley.com).

[Kohn, 1991]. In this study, we reevaluate twin resemblance in oscillatory brain activity using magnetoencephalogram (MEG) recordings. Compared to EEG which measures tangential as well as radially oriented currents, MEG is selectively sensitive to cortical sources with tangential orientation to the skull surface, i.e., the electromagnetic activity from cortical fissures [Hamalainen et al., 1993]. Nevertheless, given that both recording techniques detect synchronized neuronal currents, EEG and MEG signals can be regarded as functionally related which also has been confirmed empirically, e.g., [Kirsch et al., 2007; Nikulin et al., 2005]. In contrast to EEG however, MEG measures magnetic fields resulting from electrical brain activity which are virtually undistorted by tissue covering the brain [Okada et al., 1999; Wolters et al., 2006].

METHODS

Participants

Our sample consisted of 20 healthy right-handed MZ twin pairs (10 male, 18.8 ± 0.6 yrs; 10 female, 18.8 ± 0.4 yrs). Zygosity was based on buccal cell DNA typing. All twins provided informed consent and the study was approved by the medical ethics committee of the Vrije Universiteit medical center.

MEG Recordings

Magnetic brain activity was recorded for 10 min during eyes-closed rest, using a 151 sensor whole-head scanner with axial gradiometers (VSM Medtech, Canada). Sampling rate was 625 Hz, with low-pass filtering at 265 Hz.

Before and after each measurement head position was determined and, to correct for the influence of head position on recorded amplitudes, MEG data for each individual were extrapolated onto new data sets with the same sensor locations corresponding to an average head position across twins [de Munck et al., 2001].

Data Processing

MEG signals were processed using Fieldtrip software (F.C. Donders Centre for Cognitive Neuroimaging; <http://www.ru.nl/fcdonders/fieldtrip>). The 10 min recording sessions were partitioned into 120 epochs of 5 s. Epochs containing artifacts including excessive amplitude offsets and muscle activity were excluded after visual inspection. Eye movement and eye blink signals were removed by means of independent component analysis available in fieldtrip. For each 5 s epoch, MEG power was determined by applying a Fast Fourier Transform on the signal from each sensor, after subtracting the mean MEG amplitude and convolving the data with a Hanning taper. Smoothed powers between 1.5 and 30 Hz, with 0.2 Hz resolution, were computed by averaging the power values over epochs. Finally, power values were summed separately for the four classical frequency bands, delta (1.6–3.6 Hz), theta (4.0–7.6 Hz), alpha (8.0–12.6 Hz), and beta (13.0–25.0 Hz) and transformed to a Gaussian distribution using $\log_{10}(x)$ [Pivik et al., 1993]. Similarities of log transformed power values between MZ twins were quantified, for each recording sensor, using Pearson's linear correlation coefficient. There were no indications of sex differences, therefore correlations were determined across the entire sample of male and female twin pairs.




Figure 1.

Scalp topographies of twin correlations for \log_{10} transformed mean powers, across all 20 MZ twin pairs, for each MEG frequency band. Color coding ranges from darkblue (correlation: 0) to darkred (correlation: 1). As is immediately clear from the figure, twin resemblance in MEG power was high to very high across the scalp in the delta (1.6–3.6 Hz), theta (4.0–7.6 Hz), alpha (8.0–12.6 Hz), and beta (13.0–25.0 Hz) frequency bands.

Small map inserts below each correlation map show distributions of absolute powers (darkblue: 0 to darkred 3.2 pT²). Delta and theta bands exhibit a clear posterior/temporal power distribution. As classically observed during rest, highest power was observed in the alpha frequency range, peaking around 10 Hz, and with posterior scalp maximum. Scalp topography for the beta band was similar to alpha, except for lower power.

TABLE I. MZ twin correlations per sensor group

	RO (10)	LO (9)	RP (9)	LP (9)	RC (15)	LC (15)	RT (21)	LT (21)	RF (16)	LF (15)	Z (9)	Sensor mean
Delta	0.82 (0.59–0.93)	0.82 (0.59–0.93)	0.82 (0.59–0.93)	0.87 (0.70–0.95)	0.86 (0.67–0.94)	0.85 (0.65–0.94)	0.69 (0.36–0.87)	0.56 (0.16–0.80)	0.84 (0.63–0.94)	0.66 (0.31–0.85)	0.87 (0.70–0.95)	0.79
Theta	0.90 (0.76–0.96)	0.89 (0.74–0.96)	0.87 (0.70–0.95)	0.92 (0.81–0.97)	0.89 (0.74–0.96)	0.91 (0.78–0.96)	0.76 (0.48–0.90)	0.73 (0.43–0.89)	0.86 (0.67–0.94)	0.69 (0.36–0.87)	0.92 (0.81–0.97)	0.85
Alpha	0.94 (0.85–0.98)	0.94 (0.85–0.98)	0.89 (0.74–0.96)	0.90 (0.76–0.96)	0.84 (0.63–0.94)	0.84 (0.63–0.94)	0.85 (0.65–0.94)	0.85 (0.65–0.94)	0.86 (0.67–0.94)	0.77 (0.50–0.90)	0.95 (0.88–0.98)	0.88
Beta	0.89 (0.74–0.96)	0.91 (0.78–0.96)	0.88 (0.72–0.95)	0.94 (0.85–0.98)	0.90 (0.76–0.96)	0.89 (0.74–0.96)	0.78 (0.52–0.91)	0.81 (0.57–0.92)	0.90 (0.76–0.96)	0.80 (0.55–0.92)	0.94 (0.85–0.98)	0.88
Band mean	0.89	0.89	0.87	0.91	0.87	0.87	0.77	0.74	0.87	0.73	0.92	

Pearson's correlations (and 95% confidence intervals) between MZ twins of log10 transformed mean powers, averaged across sensors for 11 MEG sensorgroups defined according to scalp location: LO and RO: left and right occipital; LP and RP: parietal; LC and RC: central; LT and RT: temporal; LF and RF: frontal; Z: midline sensors. The number of individual sensors in each sensorgroup is indicated within brackets (one left occipital and one left frontal MEG sensor was excluded due to bad recording quality). All correlation coefficients were significant at $p < 0.001$, except for the delta band at sensorgroups LT and LF: $p \leq 0.01$.

RESULTS

MZ twin correlations of log transformed mean powers for each sensor across the scalp in Figure 1 and across sensors grouped according to scalp subdivision in Table I clearly indicate that within-pair MZ twin correlations for MEG power in all classical broad frequency bands were very high across the scalp and highly similar to those previously reported for EEG [Smit et al., 2005; Van Baal et al., 1996; van Beijsterveldt et al., 1996; Zietsch et al., 2007]. Also in agreement with EEG findings [Van Baal et al., 1996; van Beijsterveldt et al., 1996], correlations were relatively reduced for slower brain rhythms, especially for the delta band (Table I: 0.79 across sensors). This may be indicative of environmental influences on slow waves but, because reduced correlations were observed mainly at bilateral temporal and left frontal scalp locations, most likely reflects the lower signal-to-noise level at these anterior brain sites. In addition, there may be additional error variance introduced by residual biomagnetic activity from eye movement and/or eye blink signals.

DISCUSSION

Our finding of high correlations of brain activity recorded with MEG between genetically identical MZ twins, in keeping with earlier EEG studies, falsifies the idea that MZ twin resemblance for brain activity is merely a byproduct of greater MZ similarity in intervening biological tissue. The current evidence from EEG studies in twins can therefore be safely interpreted to indicate a substantial influence of genetic factors on brain activity.

ACKNOWLEDGMENT

We thank Jeroen Verbunt for technical support.

REFERENCES

- Babiloni F, Babiloni C, Carducci F, Del Gaudio M, Onorati P, Urbano A (1997): A high resolution EEG method based on the correction of the surface Laplacian estimate for the subject's variable scalp thickness. *Electroencephalogr Clin Neurophysiol* 103:486–492.
- de Munck JC, Verbunt JP, van't Ent D, van Dijk BW (2001): The use of an MEG device as 3D digitizer and motion monitoring system. *Phys Med Biol* 46:2041–2052.
- Hamalainen M, Hari R, Ilmoniemi RJ, Knuutila J, Lounasmaa OV (1993): Magnetoencephalography—Theory, instrumentation, and applications to noninvasive studies of the working human brain. *Rev Modern Phys* 65:413–497.
- Kirsch HE, Mantle M, Nagarajan SS (2007): Concordance between routine interictal magnetoencephalography and simultaneous scalp electroencephalography in a sample of patients with epilepsy. *J Clin Neurophysiol* 24:215–231.
- Kohn LAP (1991): The role of genetics in craniofacial morphology and growth. *Ann Rev Anthropol* 20:261–278.

-
- Leissner P, Lindholm LE, Petersen I (1970): Alpha amplitude dependence on skull thickness as measured by ultrasound technique. *Electroencephalogr Clin Neurophysiol* 29:392–399.
- Nikulina VV, Nikulina AV, Yamashita H, Rossi EM, Kahkonen S (2005): Effects of alcohol on spontaneous neuronal oscillations: A combined magnetoencephalography and electroencephalography study. *Prog Neuropsychopharmacol Biol Psychiatry* 29:687–693.
- Okada YC, Lahteenmaki A, Xu C (1999): Experimental analysis of distortion of magnetoencephalography signals by the skull. *Clin Neurophysiol* 110:230–238.
- Pivik RT, Broughton RJ, Coppola R, Davidson RJ, Fox N, Nuwer MR (1993): Guidelines for the recording and quantitative analysis of electroencephalographic activity in research contexts. *Psychophysiology* 30:547–558.
- Smit CM, Wright MJ, Hansell NK, Geffen GM, Martin NG (2006): Genetic variation of individual alpha frequency (IAF) and alpha power in a large adolescent twin sample. *Int J Psychophysiol* 61:235–243.
- Smit DJ, Posthuma D, Boomsma DI, Geus EJ (2005): Heritability of background EEG across the power spectrum. *Psychophysiology* 42:691–697.
- Toga AW, Thompson PM (2005): Genetics of brain structure and intelligence. *Annu Rev Neurosci* 28:1–23.
- Van Baal GC, De Geus EJ, Boomsma DI (1996): Genetic architecture of EEG power spectra in early life. *Electroencephalogr Clin Neurophysiol* 98:502–514.
- van Beijsterveldt CE, Boomsma DI (1994): Genetics of the human electroencephalogram (EEG) and event-related brain potentials (ERPs): A review. *Hum Genet* 94:319–330.
- van Beijsterveldt CE, Molenaar PC, De Geus EJ, Boomsma DI (1996): Heritability of human brain functioning as assessed by electroencephalography. *Am J Hum Genet* 58:562–573.
- Vogel F, Motulsky AG (1997): Brain physiology: Genetics of the EEG. In: Vogel F, Motulsky AG, editors. *Human Genetics: Problems and Approaches*, 3rd ed. Berlin: Springer-Verlag. pp 674–677.
- Wolters CH, Anwander A, Tricoche X, Weinstein D, Koch MA, MacLeod RS (2006): Influence of tissue conductivity anisotropy on EEG/MEG field and return current computation in a realistic head model: A simulation and visualization study using high-resolution finite element modeling. *Neuroimage* 30:813–826.
- Zietsch BP, Hansen JL, Hansell NK, Geffen GM, Martin NG, Wright MJ (2007): Common and specific genetic influences on EEG power bands delta, theta, alpha, and beta. *Biol Psychol* 75:154–164.